

Highlights of Analytical Sciences in Switzerland

Division of Analytical Sciences

A Division of the Swiss Chemical Society

Mass Spectrometric Characterization of Disulfide Bridges in Snake Venom Proteins

Miriam S. Goyder*, Marc E. Pfeifer, and Franka Kálmán

*Correspondence: Dr. M. S. Goyder, Institute of Life Technologies, University of Applied Sciences Western Switzerland, Route du Rawyl 47, CH-1950 Sion, Tel.: +41 27 606 86 40, E-mail: miriam.goyder@hevs.ch

Keywords: Disulfide bridges · Mass spectrometry · Protein analysis · Snake venom proteins


Venoms from snakes, spiders, scorpions *etc.* offer an Aladdin's cave of proteins and peptides for drug discovery. Within the venom of these species are numerous proteins and enzymes designed to kill or immobilise prey and aid digestion, including neurotoxins, cardiotoxins and muscarinic toxins. The individual proteins can be isolated (or produced recombinantly) and used for therapeutic effect.

Disulfide bridges are covalent bonds formed between the thiol groups of two cysteine residues. Many venom proteins contain a proportionally large number of such bridges, which play a fundamental role in defining their structure and specialized functionality. For many proteins, disulfide assignment can be achieved fairly simply by mass spectrometry (MS) analysis of bridged fragments following proteolytic digestion under non-reducing conditions. In venom proteins, elucidation of disulfide connectivities is often much more challenging due to the interwoven arrangement of bridges, which means that a simple proteolytic digest will not cut the protein in such a way that the connectivity can be determined unambiguously. Thus a variety of techniques are required to yield fragments that will reveal the disulfide connectivity.

At HES-SO Valais, recombinant snake venom proteins are being developed for use as new drugs. Part of the analytical strategy is to verify that disulfide bridges are correctly formed. To confirm their number, a high mass accuracy QTOF mass spectrometer is used to observe the mass difference of 1 Da between a disulfide bonded and free cysteine. The presence of disulfide isoforms (proteins with identical sequence but different




Artist's representation of mambagin-1, a pain-relieving peptide from the venom of the black mamba snake. The peptide contains four disulfide bonds in the positions Cys1-3, 2-4, 5-6, and 7-8, as shown by the grey lines. Image reproduced with permission of John Wiley and Sons, from C. J. Craik, C. I. Schroeder, *Angew. Chem. Int. Ed.* **2013**, 52, 3071.



3D model of the three-finger toxin, mambin. Disulfide bonds are shown in red, yellow, light blue and dark blue. Image of PDB ID: 2LA1 (C.H. Cheng *et al.*, *Protein Sci.* **2012**, 21, 1872) created using RCSB PDB Protein workshop (L. Moreland *et al.*, *BMC Bioinformatics* **2005**, 6, 1472).

interwoven disulfides and vicinal cysteines as are often an occurrence in venom proteins. **Verifying disulfide bonding in venom protein-based drugs is a key element in assuring Active Pharmaceutical Ingredient safety and efficacy.**

Received: December 19, 2013



MS/MS disulfide connectivity determination strategy. Xaa represents any amino acid in a hypothetical protein, which has undergone enzymatic digestion to yield disulfide bridged chains with two possible disulfide connectivities: Cys1-Cys3 or Cys2-Cys3. In this case, b2 and y3 ions from fragmentation on the first chain elucidate the connectivity. Reproduced with permission of John Wiley and Sons, from M. S. Goyder, F. Rebeaud, M. E. Pfeifer, F. Kálmán, *Exp. Rev. Proteomics* **2013**, 10, 489.

Can you show us your analytical highlight?

Please contact: Dr. Veronika R. Meyer, EMPA St.Gallen, Lerchenfeldstrasse 5, 9014 St.Gallen
Phone: +41 58 765 77 87, Fax: +41 58 765 77 62, Mail to: veronika.meyer@empa.ch